


Formacja Jura.

Taf I.



R. Fleck psr

in Lit. M. Fleck & C° à Wissamie.

Ammonites Harveyi Sowerby. M.B.







Fig. 1. *Ammonites Herveyi*, Sowerby W.B.

Fig. 2-5. *A. hereticus* Rönecke W.B.

Am. Hadolii. Somerby. W. J. A.


or Lat. J. F. Fleck & C°

J. F. Fleck sp.



Formae Jura.


Tab. IV.



H. Fleck 1852.

in Lit. J. V. Fleck & C. in Harenberg.


Terebratula concinna, Sowerby, W.B.



Pl. V. figs.

Fig. 1-5 *Terebratula trilobata* Munster W.K.Fig. 6-10 *T. varians*, Schlotheim W.B.

nella DE Puydt



R. Fleckrys:


re Lit: J.F. Fleck & C.

Terebratula biplicata, Somerby.

Fig. 1. 2. *Caryophyllia crassa*. Zel'sz. P.S.

Bockanski Z. P.S.


Fig. 3. 4.
Fig. 5. 6. *Turbinalia sinuosa*. Al. Brongnart P.P.



Formaria Jura.

Pl. VIII.






R. Płocki rys.

w lit. J. U. Pełek t. C. w Warszawie.

Fig. 1-3. *Natica compressa*, Basterot P.Z.

Fig. 3-6. *mille punctata*, Lamark P.Z.



Fig. 7-9. *glaucina*, Lamark P.Z.

Fig. 1-6. *Terebratula inconstans*. Sowerby W.B.Fig. 7-11. *tetraedra*. Sowerby W.M.B.

R. Fleck 1914


Gymnella vesicularis, Bonn 9

no. 1, top. J. F. Fleck & C. W. Warman




47
lirede

Forma cya Křídowa




B. Riedl.


Tab. VII.



n. Linn. J. T. Knob & C. de Barrois et

Cryptidea vesicularis, Brongn. O.






R. Peale type

Ammonites hyperboreus, Sowerby H. T. H.

Orthacya wezlowa.

1



"ab. Y"

2

